AD-A104 335   NAVAL SURFACE WEAPONS CENTER DAHLGREN VA   F/G 8/3
                GLOBAL OCEAN TIDES. PART V. THE DIURNAL PRINCIPAL LUNAR TIDE (O--ETC(U)
                MAY 81   E W SCHWIDERSKI
UNCLASSIFIED   NSWC/TR-81-144   NL

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER NSWC TR 81-144	2. GOVT ACCESSION NO. AD-A104	3. RECIPIENT'S CATALOG NUMBER 335
4. TITLE (and Subtitle) GLOBAL OCEAN TIDES, PART V: THE DIURNAL PRINCIPAL LUNAR TIDE (O1), ATLAS OF TIDAL CHARTS AND MAPS		5. TYPE OF REPORT & PERIOD COVERED Final
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) E. W. Schwiderski		8. CONTRACT OR GRANT NUMBER(s)
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Surface Weapons Center (K104) Dahlgren, Virginia 22448		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61152N/R0000-1 ZR000-01-01/1K01AA
11. CONTROLLING OFFICE NAME AND ADDRESS Chief of Naval Material Department of the Navy Washington, DC 20360		12. REPORT DATE May 1981
		13. NUMBER OF PAGES 85
14. MONITORING AGENCY NAME & ADDRESS(if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE

16. DISTRIBUTION STATEMENT (of this Report)

Approved for public release; distribution unlimited.

17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)

18. SUPPLEMENTARY NOTES

19. KEY WORDS (Continue on reverse side if necessary and identify by block number)

Ocean Tides and Currents
Numerical Modeling
Tidal Charts

20. ABSTRACT (Continue on reverse side if necessary and identify by block number)

In Part I (Schwiderski, 1978a) of this report, a unique hydrodynamical interpolation technique was introduced, extensively tested, and evaluated in order to compute partial global ocean tides in great detail and with a high degree of accuracy. This novel method has been applied to construct the diurnal principal lunar (O_1) ocean tide with a relative accuracy of better than 5 cm anywhere in the open oceans. The resulting tidal amplitudes and phases are tabulated on a

(see back)

DD FORM 1473 EDITION OF 1 NOV 65 IS OBSOLETE
1 JAN 73
S/N 0102-LF-014-6601

(20)

$1° \times 1°$ grid system in an atlas of $42° \times 71°$ overlapping charts covering the whole oceanic globe. A corresponding atlas of global corange and cotidal maps is included to provide the reader with a quick general overview of the major tidal phenomena. The specifying hydrodynamical parameters of the model are listed along with quoted sources of empirical tide data, and significant tidal features are explained and discussed. The diurnal O_1 ocean tide is found to resemble closely the diurnal K_1 tide and qualitatively also the semidiurnal S_2 and M_2 tides which were presented in Parts IV, III, and II of this report, respectively.

FOREWORD

In Part I of this report (Schwiderski, 1978a), a combined hydrodynamical-empirical method was introduced to compute numerically harmonic partial tides in the world oceans with an accuracy of better than 5 cm, which is needed in various military and civil applications of today. In this report, the computed diurnal principal lunar tide (O_1) is displayed in an atlas of tabulated tidal charts and plotted corange and cotidal maps.

The date of completion was May 20, 1981.

Released by

R. T. RYLAND, JR., Head
Strategic Systems Department

*National Ocean Survey (NOS)
 National Oceanographic and Atmospheric Administration (NOAA)

iii

CONTENTS

ABSTRACT

In Part I (Schwiderski, 1978a) of this report, a unique hydrodynamical interpolation technique was introduced, extensively tested, and evaluated in order to compute partial global ocean tides in great detail and with a high degree of accuracy. This novel method has been applied to construct the diurnal principal lunar (O_1) ocean tide with a relative accuracy of better than 5 cm anywhere in the open oceans. The resulting tidal amplitudes and phases are tabulated on a $1° \times 1°$ grid system in an atlas of $42° \times 71°$ overlapping charts covering the whole oceanic globe. A corresponding atlas of global corange and cotidal maps is included to provide the reader with a quick general overview of the major tidal phenomena. The specifying hydrodynamical parameters of the model are listed along with quoted sources of empirical tide data, and significant tidal features are explained and discussed. The diurnal O_1 ocean tide is found to resemble closely the diurnal K_1 tide and qualitatively also the semidiurnal S_2 and M_2 tides which were presented in Parts IV, III, and II of this report, respectively.

I. INTRODUCTION

Part 1 of this report (Schwiderski, 1978a) introduced a unique combination of hydrodynamical and empirical methods to model detailed ocean tides with a relative component accuracy of better than 5 cm anywhere in the open oceans. This enormous accuracy is well above minimum requirements set by, for instance, the National Aeronautics and Space Administration (NASA) and the Department of Defense (DoD) to map the geoid at sea by satellite altimetry to within 10 cm. The following features of this unique hydrodynamical interpolation model made the achievement of this accuracy possible.

a. A spherically graded $1° \times 1°$ grid system is set up in connection with a corresponding $1° \times 1°$ bathymetry to assure a sufficient resolution of all important tidal phenomena.

b. The bathymetry of the gridwise, simply connected ocean basin is hydrodynamically defined (Schwiderski, 1978c) by appropriate modifications of earlier realistic depth data collections. The hydrodynamical redefinition was needed in order to model the well-known strong distortion and retardation effects of shallow continental shelves, narrow ocean ridges or island chains, and other significant bottom irregularities.

c. The Boussinesq substitution of the turbulent Reynolds stresses is applied in the form of eddy dissipation with a novel physically meaningful eddy viscosity that depends linearly on the lateral grid-cell area and, hence, directly on the ocean depth.

d. The linear law of bottom friction is introduced with a bottom-friction coefficient depending linearly on the bottom grid-cell area which is independent of the ocean depth. In boundary cells, the otherwise constant friction coefficient is subjected to an indirect cellwise adjustment in order to permit a consistent hydrodynamical interpolation (see h., below) of empirical tide data known from tide gauge stations at continental shores, islands, or other shallow-ocean bottom irregularities.

e. The effects of the terrestrial tide and the oceanic tidal load are included as simple second-order approximations in the sense of Love and Accad and Pekeris (1978).

f. The Hansen-Zahel (Zahel, 1970 and 1977; Estes, 1977) finite differencing technique is modified by a new differencing scheme in time which improved decay, dispersion, and stability characteristics of the numerical procedure and facilitates the simple indirect adjustment of the bottom-friction coefficient in the hydrodynamical interpolation technique (see d. and h.).

g. At land-ocean cell walls, the conditions of no-flow across and free-slip along the boundaries are enforced. The no-flow condition is subsequently relaxed by allowing controlled periodic inflows and outflows over the mathematically assumed boundaries. This allowance redefines indirectly more realistic shorelines in order to further improve the consistency of the hydrodynamical interpolation of empirical data (see d. and h.).

1

h. A unique hydrodynamical interpolation technique is introduced which incorporates into the theoretical model empirical tidal constants collected from over 2 000 tide-gauge stations around the world in a hydrodynamically consistent fashion (see d., f., and g., above).

i. A new higher order approximation of Arctic Ocean tides is used, that is described in Schwiderski (1981c).

With these features, the new model was successfully applied to chart the semidiurnal principal lunar (M_2) ocean tide with the desired accuracy. The technique and accuracy of the model were extensively described and discussed in Part I of this report as well as in subsequent journal publications and symposia presentations by the author (Schwiderski 1978a, b; 1979a, b, c, d, e; and 1980).

The same hydrodynamical interpolation technique has been applied to chart the diurnal principal lunar (O_1) ocean tide with the same relative accuracy as M_2. Again, it must be emphasized that the enormous accuracy achieved over all open ocean regions diminishes somewhat near coastal areas where known empirical data are marginal in quantity and/or quality.

A complete listing of all sources of empirical ocean tide data, which were interpolated into the O_1 tidal charts, is presented in Appendix A. In the meantime, Section 2 of this report lists the significant hydrodynamical input parameters that specified the constructed O_1 ocean tide. The major features of the global O_1 tide are discussed in Section 3. A complete numerical display is presented in Appendix A where all tidal amplitudes and phases are gridwise tabulated in map-like charts. Corange (equi-amplitude) and cotidal (equi-phase) maps of the O_1 ocean tide are plotted in Appendix B.

2. O_1 OCEAN-TIDE PARAMETERS

The astronomical diurnal principal lunar (O_1) equilibrium tide η (or tide-generating potential $G\eta$; see Schwiderski, 1978a) at the geographical point (λ, ϕ) and instant (Y, D, t) is determined by

$$\eta = K \sin 2 \phi \cos(\sigma t + X + \lambda) \tag{1}$$

where

G = 9.81 m/sec^2 earth gravity acceleration
λ = longitude (east in rad)
ϕ = latitude (north in rad)
Y ($\geqslant$ 1975) = year number
D = day number of year Y(D = 1 for January 1)
t = universal standard time of day D (in sec)
K = 0.100 574 m = O_1 equilibrium tide amplitude
σ = 0.67598 $\cdot$ 10^{-4} sec^{-1} = O_1 tide frequency
X = π (h_0 - 2 s_0 - 90)/180 = O_1 astronomical argument (in rad)
h_0 $\begin{cases} = 279.696\ 68 + 36\ 000.768\ 930\ 485T + 3.03 \cdot 10^{-4}\ T^2 \\ = \text{mean longitude of the sun relative to Greenwich midnight of day } D \text{ (in deg)} \end{cases}$
s_0 $\begin{cases} = 270.434\ 358 + 481\ 267.883\ 141\ 37T - 0.001\ 133T^2 + 1.9 \cdot 10^{-6}\ T^3 \\ = \text{mean longitude of the moon relative to Greenwich midnight of day } D \text{ (in deg)} \end{cases}$
T = [27 392.500 528 + 1.000 000 035 6$\bar{D}$]/36 525
$\bar{D}$ = D + 365 (Y - 1975) + Int [(Y - 1973)/4]
Int[x] = integral part of x

The corresponding instantaneous ocean partial tide (Schwiderski, 1978a) is determined by

$$\zeta = \xi \cos (\sigma t + X - \delta), \tag{2}$$

where the local harmonic constants

$\xi = \xi(\lambda, \phi) = O_1$ ocean tide amplitude (in m)

and

$\delta = \delta(\lambda, \phi) = O_1$ ocean tide Greenwich phase (in rad)

must be determined, say, by linear interpolation in the tidal charts of Appendix A.

A simple second-order approximation in the sense of Love and Accad and Pekeris (see Part I, Schwiderski, 1978a, 1979c, and 1980; and Accad and Pekeris, 1978) yields

3

$$\zeta' \approx 0.612\eta \text{ and } \zeta^{eo} \approx -0.0667\zeta,$$ (3)

i.e., the corresponding terrestrial tide ζ^e and the earth dip ζ^{eo} (yielding) under the oceanic tidal load ζ, respectively. A more elaborate and probably slightly more accurate earth dip ζ^{eo} may be computed by using Farrell's Green function (see Farrell, 1972 and 1973; and Schwiderski, 1980). In linear superposition, one finds the corresponding instantaneous geocentric partial O_1 tide:

$$\zeta^c = \zeta + \zeta^c + \zeta^{eo}.$$ (4)

A detailed description of the hydrodynamical-empirical model to compute the ocean tidal amplitudes ξ and phases δ (listed in Appendix A) was given in Schwiderski (1978a, 1979c, d, and 1980). In particular, all model input parameters such as the dimensionless eddy coefficient ϵ (Eq's. 103 and 123), the bottom-friction parameter b (Eq's. 4a and b), and the differencing parameters κ and $\bar{\kappa}$ (Eq's. 64 and 72) were all specified in Schwiderski (1978a) (referenced equations). These parameters were determined for M_2 by extensive trial-and-error computations and remained unchanged for the construction of O_1.

In the computation of the O_1 tide model, the following mode-dependent parameters were used (see referenced equations in Schwiderski, 1978a):

a. The time step Δt (Eq's. 64, 123):

$$\Delta t = 193.6443 \text{ sec}$$ (5)

b. The hydrodynamical interpolation control limits, k_1, k_2, and k_3 (Eq's. 88, 89, 94, 97, and 99)

$$k_1 = 0.025, k_2 = 0.040, k_3 = 0.5.$$ (6)

It may be noted that the input parameters k_1 and k_2 of Equations 6 are the same as for the diurnal K_1 component, but different from those values used for the semidiurnal S_2 and M_2 species (see Parts IV, III, and II).

4

3. O_1 OCEAN-TIDE FEATURES

The entire constructed O_1 ocean tide is gridwise displayed in map-like amplitude and phase tables in Appendix A. The 42° x 71° charts cover the whole globe north of colatitude 169° (Antarctica) in three zones: a northern zone N from 0° to 71° colatitude, a middle zone M from 48° to 118° colatitude, and a southern zone S from 98° to 168° colatitude. The overlapping geographical areas of the tidal charts have been chosen to provide a worldwide coverage for special applications and to allow the reader to scan the large amplitude and phase charts together in order to evaluate their quality and visualize the important tidal features. In addition, a generally superficial overview of some tidal features can be recognized by inspecting the more schematically plotted corange and cotidal maps provided in Appendix B.

For an easy evaluation of the tidal charts in Appendix A, all hydrodynamically interpolated empirical tidal amplitudes and phases have been visibly marked by subbars for all shore data and subbrackets for all near-shore deep-sea input constants. Furthermore, the charts display the approximate locations of distant off-shore deep-sea stations by subtildes under the computed amplitude and phase data. The corresponding empirical data, which were excluded from hydrodynamical interpolation (see Sect. 1 and Schwiderski, 1978a, 1979d, and 1980), are listed and compared with the modeled data in Tables 1, 2, and 3. Finally, the approximate geographical locations of the important amphidromic points of zero amplitudes are marked by a circled $\otimes$.

The tidal charts and maps permit the viewer to follow the tidal waves, that is the high water fronts (crests), in forward (or backward) direction, for instance, on their rotation around the amphidromic points. In the tidal phase charts of Appendix A, it is best to start from the prominently visible 0° = 360° or 100° cotidal lines. Since the Greenwich phases specify the time lags (in degrees: 15° ≈ 1 hour) of the tidal crests relative to the cresting time of the corresponding equilibrium tide along Greenwich meridian, one gathers a vivid impression of the significant global and local tidal phenomena.

By following the tidal waves on their periodic rotations, one finds these waves passing through the specially marked stations in empirically correct time and with the correct height. In fact, all over the globe over 2 000 tidal phases and 2 000 amplitudes are coherently integrated. This is particularly impressive for the charts of the Pacific Ocean, where the empirical data from so many clustered and scattered island stations fit smoothly into the surrounding computed tides. From the smoothness features of erratically interpolated tidal data (see Parts I and II), one concludes that this result is not an artifact of the interpolation applied but constitutes a vivid manifestation of the excellent compatibility of both the empirical and hydrodynamical procedures combined.

On the basis of this observation, it can again (see Schwiderski, 1978a, b; 1979a, b, d, e; 1980, and 1981a, b) be estimated that the O_1 tidal charts permit a tide prediction with a uniform accuracy relative to M_2 of better than 5 cm anywhere in the open oceans. Naturally, near rough ocean basin reliefs (e.g., Arctic and Antarctic shores), where empirical tide (and depth) data are marginal in quality and quantity, a somewhat lesser accuracy must be expected. The estimated

5

accuracy of the computed O_1 tide is, of course, fully validated by all 32 empirical tide data from distant off-shore deep-sea tide gauge stations, which are listed along with the computed data in Tables 1, 2, and 3. The differences (not necessarily errors) range from 0 to 1 cm in amplitudes and 0° to 11° (44 minutes) in phases and thus verify the estimated prediction accuracy. In this connection one may recall the accuracy evaluation of the deep-sea empirical data presented in Part IV of this report.

Table 1. North Atlantic Ocean Deep-Sea Empirical and Modeled O_1 Tides

LONG W	LAT N	EMP ξ	MOD ξ	Δξ	EMP δ	MOD δ	Δδ	IAPSO NR	SOURCES
13°51'	58°16'	7	6	−1	16	13	−3	1.1.37	C
24°43'	62°50'	6	6	0	75	64	−11	1.1.29	C
28°46'	60°12'	5	5	0	66	66	0	1.1.30	C
29°58'	57°01'	5	4	−1	66	58	−8	1.1.31	C
30°10'	53°39'	3	3	0	57	51	−6	1.1.32	C
25°06'	53°31'	4	4	0	19	25	+6	1.1.33	C
20°00'	53°39'	5	5	0	9	9	0	1.1.34	C
28°11'	48°45'	3	2	−1	26	22	−4	1.1.38	C
28°09'	45°21'	2	2	0	10	8	−2	1.1.39	C
27°57'	41°25'	2	2	0	342	343	+1	1.1.40	C
20°05'	37°09'	4	3	−1	318	319	+1	1.1.41	C
14°15'	36°41'	6	5	−1	316	314	−2	1.1.42	C
75°38'	32°42'	8	7	−1	192	191	−1	1.2. 3	C.M
76°25'	30°26'	7	7	0	194	196	+2	1.2.11	C, P
76°48'	28°27'	7	7	0	196	198	+2	1.2.15	C
76°47'	28°01'	7	7	0	202	198	−4	1.2.14	C
67°32'	28°14'	6	5	−1	197	200	+3	1.2. 5	C, Z
69°45'	28°08'	6	6	0	198	199	+1	1.2. 4	C, Z
69°40'	27°59'	7	6	−1	201	201	0	1.2. 8	C, Z
69°40'	27°58'	6	6	0	196	201	+5	1.2. 7	C, Z
69°20'	26°28'	6	6	0	200	204	+4	1.2.10	C, Z
69°19'	26°27.	6	6	0	199	204	+5	1.2. 9	C, Z

ξ   Amplitudes (cm)
δ = Greenwich Phases (deg)
IAPSO = Int. Assoc. for the Phys. Sci. of the Oceans
C = Cartwright et al. (1979)
M = Mofjeld (1975)
P = Pearson (1975)
Z = Zetler et al. (1975)

Table 2. Northeastern Pacific Ocean Deep-Sea Empirical and Modeled O_1 Tides

LONG W	LAT N	EMP ξ	MOD ξ	Δξ	EMP δ	MOD δ	Δδ	IAPSO NR	SOURCES
144°22'	56°08'	27	28	+1	250	253	+3	2.1.17	C
135°38'	53°19'	28	28	0	244	244	0	2.1.16	C
132°47'	49°35'	26	26	0	231	235	+4	2.1.15	C
145°00'	34°00'		15	–	–	227	–	...	--
145°00'	34°00'		15	–	–	227	–	--	--
124°26'	27°45'	18	17	–1	199	199	0	2.1.13	C, M
129°01'	24°47'	16	15	–1	201	204	+3	2.1.10	C, M

ξ = Amplitudes (cm)
δ = Greenwich Phases (deg)
IAPSO = Int. Assoc. for the Phys. Sci. of the Oceans
C = Cartwright et al. (1979)
M = Munk et al. (1970)

Table 3. Southeast Indian Ocean Deep-Sea Empirical and Modeled O_1 Tides

LONG E	LAT S	EMP ξ	MOD ξ	Δξ	EMP δ	MOD δ	Δδ	IAPSO NR	SOURCES
132°01'	37°01'	14	13	–1	218	219	+1	4.1. 1	C, IS
132°09'	50°02'	12	11	–1	220	221	+1	4.1. 2	C, IS
132°07'	60°01'	15	16	+1	215	214	–1	4.1. 3	C, IS

ξ = Amplitudes (cm)
δ = Greenwich Phases (deg)
IAPSO = Int. Assoc. for the Phys. Sci. of the Oceans
C = Cartwright et al. (1979)
IS = Irish and Snodgrass (1972)

From the tidal charts and maps in Appendixes A and B, one concludes that the rotating tidal waves of the diurnal O_1 tide resemble closely those of the diurnal K_1 tide. There is also a qualitative similarity to the semidiurnal S_2 and M_2 tides. However, the distribution of the amphidromic systems between the diurnal and semidiurnal species varies considerably (compare Parts II, III, and IV). Also, as was mentioned for K_1 the computed and empirical distortions and retardations of the O_1 waves by boundary and bottom irregularities are generally significantly subdued when compared to the rougher semidiurnal tides as S_2 and M_2.

4. CONCLUSIONS

The hydrodynamical interpolation technique has been applied to construct the diurnal principal lunar tide (O_1) with a relative accuracy of better than 5 cm anywhere in the open oceans. The constructed tide is displayed by tabulated charts in Appendix A and by corange and cotidal maps in Appendix B. The major features of the O_1 tide are discussed in Section 3. A comparison with the earlier computed diurnal K_1 tide reveals close similarities. However, only qualitative similarities exist between the diurnal and semidiurnal species as M_2 and S_2.

REFERENCES

1. Accad, Y. and Pekeris, C. L., 1978. "Solution of the Tidal Equations for the M_2 and S_2 Tides in the World Oceans from a Knowledge of the Tidal Potential Alone," *Phil. Trans. Roy. Soc., London, Ser. A*, 290, p. 235.

2. British Admiralty Tide Tables, 1977. Vols. 1, 2, and 3.

3. Cartwright, D. E., Zetler, B. D., and Hamon, B. V, 1979. *Pelagic Tidal Constants*, IAPSO Publication Scientifique No. 30.

4. Defant, A., 1961. *Physical Oceanography, Vol. II*, Pergamon Press, New York.

5. Estes, R. H., 1977. *A Computer Software System for the Generation of Global Ocean Tides Including Self-Gravitation and Crustal Loading Effects*, Goddard Space Flight Center, TR-X-920-77-82, Greenbelt, Maryland.

6. Farrell, W. E., 1972. "Deformation of the Earth by Surface Loads," *Rev. Geophys. Space Phys.*, 10, p. 261.

7. Farrell, W. E., 1973. "Earth Tides, Ocean Tides and Tidal Loading," *Phil. Trans. Roy. Soc., London, Ser. A*, 274, p. 253.

8. Harris, R. A. 1904. *Manual of Tides, Part IV b, Report of the Superintendent*, U.S. Coast and Geodetic Survey, p. 313.

9. International Hydrographic Bureau, 1978. *Tides, Harmonic Constants*, Computer Tape, Monaco.

10. Irish, J. D., Munk, W. H., and Snodgrass, F. E., 1971. "M_2 Amphidrome in the Northeast Pacific." *Geophys. Fluid Dyn.*, 2, p. 355.

11. Irish, J. D. and Snodgrass, F. E., 1972. "Australian-Antarctic Tides," *Antarctic Res. Ser., Vol. 19: Antarctic Oceanology II: The Australian-New Zealand Sector*, edited by D. E. Hayes, AGU, p. 101.

12. Luther, D. S. and Wunsh, C., 1975. "Tidal Charts of the Central Pacific Ocean," *J. Phys. Oce.*, 5, p. 227.

13. Miyazaki, M., Juronuma, S., and Inoue, T., 1967. "Tidal Constants Along the Coast of Japan," *Oceanogr. Mag.*, 19, p. 13.

14. Motjeld, H. O., 1975. *Empirical Model for Tides in the Western North Atlantic Ocean*, NOAA, TR-ERL 340-AOML 19, Boulder, Colorado.

15. Munk, W. H., Snodgrass, F. E., and Wimbush, M., 1970. "Tides Offshore: Transition from California Coastal to Deep-Sea Waters," *Geophys. Fluid Dyn.*, 1, p. 161.

16. National Ocean Survey, 1942. *Tidal Harmonic Constants*, U.S. Coast and Geodetic Survey, Washington, D.C.

17. Nowroozi, A. A., 1972. "Long-Term Measurements of Pelagic Tidal Height off the Coast of Northern California," *J. Geophys. Res.*, 77, p. 434.

18. Nowroozi, A. A., Kuo, J. T., and Ewing, M., 1969. "Solid Earth and Oceanic Tides Recorded on the Ocean Floor of the Coast of Northern California," *J. Geophys. Res.*, 24, p. 605.

19. Pearson, C. A., 1975. *Deep-Sea Tide Observations off the Southeastern United States*, NOAA T. Memo. No. 17, Rockville, Maryland.

20. Pugh, D. 1979. "Sea Levels at Aldabra Atoll, Mombasa and Mahé, Western Equatorial Indian Ocean, Related to Tides, Meteorology and Ocean Circulation," *Deep-Sea Research*, 26A, p. 237.

21. Schwiderski, E. W. 1978a. *Global Ocean Tides, Part I: A Detailed Hydrodynamical Interpolation Model*, NSWC/DL TR-3866.

22. Schwiderski, E. W., 1978b. "A Detailed Hydrodynamical Interpolation Model of Worldwide Ocean Tides," presented at the Int. Symp. on Interaction of Marine Geodesy and Ocean Dynamics, Miami, Florida, October 10-15.

23. Schwiderski, E. W., 1978c. *Hydrodynamically Defined Ocean Bathymetry*, NSWC/DL TR-3888.

24. Schwiderski, E. W., 1979a. "NSWC Ocean Tide Program," presented at the NASA SEASAT ALT/POD Calibration Workshop, Austin, Texas, June 11-15.

25. Schwiderski, E. W., 1979b. "Detailed Ocean Tide Models of (N_2, M_2, S_2, K_2) and (K_1, P_1, O_1, Q_1) Including an Atlas of Tidal Charts and Maps," presented at the XVIIth General Assembly of the International Union of Geodesy and Geophysics in Canberra, Australia, December 2-15.

26. Schwiderski, E. W., 1979c. "Ocean Tides, Part I: Global Tidal Equations," *Marine Geodesy*, 3, p. 161.

27. Schwiderski, E. W., 1979d. "Ocean Tides, Part II: A Hydrodynamical Interpolation Model," *Marine Geodesy*, 3, p. 219.

28. Schwiderski, E. W., 1979e. *Global Ocean Tides, Part II: The Semidiurnal Principal Lunar Tide (M_2), Atlas of Tidal Charts and Maps*, NSWC TR 79-414.

29. Schwiderski, E. W., 1980. "On Charting Global Ocean Tides," *Reviews of Geophysics and Space Physics*, 18, No. 1.

30. Schwiderski, E. W., 1981a. *Global Ocean Tides, Part III: The Semidiurnal Principal Solar Tide (S_2), Atlas of Tidal Charts and Maps*, NSWC TR 81-122.

31. Schwiderski, E. W., 1981b. *Global Ocean Tides, Part IV: The Diurnal Luni-Solar Declination Tide (K_1), Atlas of Tidal Charts and Maps*, NSWC TR 81-142.

32. Schwiderski, E. W., 1981c. *Exact Expansions of Arctic Ocean Tides*, Naval Surface Weapons Center Technical Report in preparation.

33. Zahel, W., 1970. "Die Reproduktion Gezeitenbedingter Bewegungsvorgange im Weltozean Mittels des Hydrodynamisch-Numerischen Verfahrens," *Mitteilungen des Inst. f. Meereskunde der Univ. Hamburg, XVII*.

34. Zahel, W., 1977. "The Influence of Solid Earth Deformations on Semidiurnal and Diurnal Oceanic Tides," *Proc. IRIA Int. Colloq. on Numerical Methods of Science and Technical Computation*, Springer, Berlin.

35. Zetler, B. D., Munk, W. H., Mofjeld, H. O., Brown, W., and Dormer, F., 1975. "MODE Tides," *J. Phys. Oceanogr.*, 5, p. 430.

APPENDIX A

ATLAS OF $1° \times 1°$ O_1 OCEAN TIDE AMPLITUDE
AND PHASE CHARTS FOR $42° \times 71°$ AREAS

APPENDIX A

ATLAS OF 1° x 1° O₁ OCEAN TIDE AMPLITUDE AND PHASE CHARTS FOR 42° x 71° AREAS

1. **GUIDE TO TIDAL CHARTS**

M	$= m$: Longitude Number
N	$= n$: Colatitude Number
λ_m	$= (m - 0.5)°$: Geographical Longitude East
θ_n	$= (n - 0.5)°$: Geographical Colatitude
$\xi_{m,n}$	$= \xi(\lambda_m, \theta_n)$: Amplitude (in cm)
$\delta_{m,n}$	$= \delta(\lambda_m, \theta_n)$: Greenwich Phases (in deg.; $15° \approx 1$ h)
⊗	= Amphidromic Points
...	= Subbars Mark Empirical Input Data at Shore Stations
⊔	= Subbrackets Mark Empirical Input Data at Near-Shore Deep-Sea Stations
∼	= Subtildes Mark Approximately Distant Offshore Deep-Sea Stations with Excluded Empirical Tide Data Listed in Tables 1, 2, and 3

2. **SOURCES OF EMPIRICAL TIDE DATA**

Publications:

National Ocean Survey (1942), British Admiralty (1977), International Hydrographic Bureau (1978), Defant (1961), Miyazaki et al. (1967), Nowroozi et al. (1969), Munk et al. (1970), Zahel (1970), Irish et al. (1971), Irish and Snodgrass (1972), Nowroozi (1972), Luther and Wunsh (1975), Mofjeld (1975), Pearson (1975), Zetler et al. (1975), Cartwright et al. (1979), and Pugh (1979).

Private Communications:

D. C. Simpson (1977), National Ocean Survey, Rockville, Maryland; S. K. Gill and D. L. Porter (1978), National Ocean Survey, Rockville, Maryland; K. Wyrtki (1978), University of Hawaii, Honolulu, Hawaii, and D. E. Cartwright and D. Pugh (1978), Institute of Oceanographic Sciences, Bidston Observatory, U.K.

EUROPEAN USSR

SPITZBERGEN

NORWAY

WESTERN EUROPE

BRITISH ISLES

TABLE 1 N: 1° x 1° O₁ OCEAN TIDE GREENWICH PHASES δ (DEG)

EUROPEAN USSR

SPITSBERGEN

NORWAY

WESTERN EUROPE

BRITISH ISLES

TABLE 2№ 1° x 1° O$_1$ OCEAN TIDE AMPLITUDES ξ (CM)

LAND

NOVA ZEMLYA

JOSEF

FRANZ

CENTRAL USSR

WESTERN INDIA

PAKISTAN

IRAN

ARABIA

TABLE 2N: 1° x 1° O_1 OCEAN TIDE GREENWICH PHASES δ (DEG)

FRANZ

JOSEF

LAND

NOVA ZEMLYA

CENTRAL USSR

IRAN

ARABIA

PAKISTAN

WESTERN INDIA

TABLE 3N: 1° x 1° O_1 OCEAN TIDE AMPLITUDES ξ (CM)

SIBERIAN USSR

SEVERNAYA ZEMLYA

EASTERN INDIA

BANGLADESH

SOUTHERN CHINA

HAINAN

TABLE 3N: 1° x 1° O₁ OCEAN TIDE GREENWICH PHASES δ (DEG)

M/N	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100	101	102	103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120
1	280	280	280	280	280	280	279	279	279	279	279	278	278	278	278	278	278	277	277	277	277	277	276	276	276	276	276	276	275	275	275	275	275	275	274	274	274	274	274	274	274	273
2	276	276	275	275	274	274	273	273	272	272	271	271	270	270	269	269	268	268	268	267	261	264	266	265	265	265	264	264	263	263	263	262	262	261	261	261	260	260	260	259	259	259
3	272	272	272	271	270	269	268	267	267	266	265	265	264	263	263	262	262	261	261	260	259	259	258	258	257	257	256	256	255	255	254	254	253	253	252	252	251	251	250	250	249	249
4	270	269	268	267	266	266	265	264	263	262	261	261	260	259	258	258	257	256	254	255	254	254	253	252	252	251	250	250	249	249	248	248	247	246	246	245	245	244	244	243	243	242
5	263	262	261	261	259	259	258	257	256	255	255	254	253	253	252	251	251	250	249	248	248	248	247	246	244	245	245	244	244	243	243	242	242	241	241	240	239	239	238	238	238	237
6	255	254	254	253	252	251	250	250	249	248	248	247	247	246	246	245	245	244	244	243	242	242	241	241	240	240	240	239	239	238	238	237	237	236	236	235	235	234	234	234	233	233
7	247	246	246	245	245	244	244	243	243	242	242	242	242	241	241	240	240	240	239	238	237	237	237	236	236	236	235	235	234	234	234	233	233	232	232	232	231	231	231	230	230	230
8	239	239	239	239	238	238	238	238	238	237	237	237	237	236	236	236	235	235	234	234	234	233	233	232	232	232	232	231	231	231	230	230	230	230	229	228	229	229	228	228	228	227
9	230	230	230	230	231	231	232	232	233	233	233	233	232	232	232	231	231	231	230	230	230	230	231	231	231	230	230	230	230	229	229	228	228	228	228	227	227	227	227	228	228	227
10	237	237	237	236	236	235	235	234	233	233	233	233	232	229	267	271	273	273	271	270	269	268	228	227	229	229	226	226	226	226	226	225	225	225	225	224	224	224	224	224	224	224
11	235	235	235	236	236	235	235	234	236	231	230	231	231	256	261	264	261	264	267	271	217	227	228	226	222	221	221	221	221	221	221	221	222	225	222	222	222	222	223	223	223	223
12	236	237	237	236	238	238	240	242	245	248	251	252	254	263	268	268	269	270	271	274	274	227	227	229	229	216	216	215	214	213	212	212	213	213	214	215	215	216	217	219	220	221
13	242	242	243	245	246	247	249	250	251	254	257	260	263																							212	214	216	217	219	220	221
14	241	242	244	244	247	247	250	251	254	240																					213											
15	225	225	225	225	221	218																																				

EASTERN INDIA

BANGLADESH

SIBERIAN USSR

SEVERNAYA
ZEMLYA

SOUTHERN CHINA

HAINAN

TABLE 4N: 1° x 1° O_1 OCEAN TIDE AMPLITUDES ξ (CM)

EASTERN CHINA

NEW SIBERIA

EASTERN SIBERIAN USSR

KAMCHATKA

SEA OF JAPAN

SOUTHEASTERN JAPAN

NANPOS

GULF OF CHIHLI

KOREA

TAIWAN

TABLE 4N: 1° × 1° O₁ OCEAN TIDE GREENWICH PHASES δ (DEG)

EASTERN CHINA

GULF OF CHIHLI

KOREA

SEA OF JAPAN

SOUTHEASTERN JAPAN

NEW SIBERIA

EASTERN SIBERIAN USSR

KAMCHATKA

TAIWAN

NANPOS

TABLE 5N: 1° x 1° O₁ OCEAN TIDE AMPLITUDES ξ (CM)

ALASKA USA

EASTERN SIBERIAN USSR

WRANGEL

KAMCHATKA

TABLE 5N: 1° x 1° O₁ OCEAN TIDE GREENWICH PHASES δ (DEG)

EASTERN SIBERIAN USSR

ALASKA USA

KAMCHATKA

WRANGEL

TIANS

ALASKA

USA

NORTHWESTERN CANADA

WESTERN USA

QUEEN CHARLOTTE

Q. ELIZABETH I

FRANKLIN DISTRICT

TABLE 6N: 1° x 1° O_1 OCEAN TIDE GREENWICH PHASES δ (DEG)

ALASKA

USA

NORTHWESTERN CANADA

WESTERN USA

QUEEN CHARLOTTE

Q. ELIZABETH I.

FRANKLIN DISTRICT

TABLE 7N: 1° x 1° O$_1$ OCEAN TIDE AMPLITUDES ξ (CM)

QUEEN ELIZABETH ISLANDS

NORTH CENTRAL CANADA

CENTRAL USA

FRANKLIN DISTRICT

MEXICO

GULF OF CALIFORNIA

FLORIDA

CUBA

TABLE 7N: 1° x 1° O₁ OCEAN TIDE GREENWICH PHASES δ (DEG)

λN	239	240	241	242	243	244	245	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280
1	276	279	279	279	279	280	280	280	280	281	281	281	281	282	282	282	283	283	283	283	284	284	284	284	285	285	285	286	286	286	287	287	287	288	288	288	289	289	289	289	290	290
2	266	266	267	268	269	270	270	271	272	273	274	275	276	277	278	279	280	281	282	286	287	288	289	290	292	293	294	295	297	302	303	304	305	306	307	308
3	249	250	251	252	253	254	255	258	259	261	263	265	267	268	271	273	276	278	281	284	286	289	291	294	297	300	303	305	308	311	313	315	317	319	321	323	325	327	329	330
4	230	231	232	233	234	235	236	238	240	243	246	247	250	252	256	262	268	274	280	286	291	297	303	308	313	318	323	328	333	338	341	343	344	345	347	348	350	351	352	353
5	274	278	282	286	290	294	297	301	309	313	317	321	325	329	331	333	336	338	341	343	345	347	348	349	350	354	355	355	356	357	357
6	312	316	311	326	330	334	335	338	341	343	346	347	349	350	351	352	353	354	355	355	356	357
7	325	328	331	333	336	339	341	342	344	345	347	348	349	350	351	352	353	354	355	356	357	358	359
8	328	331	333	335	340	342	344	345	347	348	349	350	351	352	353	353	355	356	357	358	359	359
9	321	325	328	332	334	336	338	340	342	343	345	347	348	350	351	352	353	353	354	355	356	357	358	359	356
10	310	314	318	323	326	328	330	333	335	338	339	340	341	342	344	346	347	348	349	350	351	251	251
11	296	302	307	312	315	318	320	322	324	325	327	327	331	335	337	346	355	360	1	2	3
12	282	288	295	301	306	311	315	320	322	324	324	324	327	331	335	339	342	346	350	350	3
13	269	276	283	290	297	304	311	315	320	323	324	325	327

FRANKLIN DISTRICT

| 15 | 289 | 303 |
| 16 | 286 | 311 |

QUEEN ELIZABETH ISLANDS

| 20 | 200 | 201 | 202 | 203 |
| 21 | | 204 | 204 |

NORTH CENTRAL CANADA

CENTRAL USA

MEXICO

GULF OF CALIFORNIA

(Numeric data blocks for rows 56–71 in the Mexico / Gulf of California region)

FLORIDA

CUBA

TABLE 8N: 1° x 1° O_1 OCEAN TIDE AMPLITUDES ξ (CM)

GREENLAND

QUEEN ELIZABETH ISLANDS

BAFFIN ISLAND

EASTERN CANADA

NEWFOUNDLAND

NOVA SCOTIA

EASTERN USA

LONG ISLAND

HISPANIOLA

BAHAMAS

CUBA

FLORIDA

TABLE 8N: 1° x 1° O₁ OCEAN TIDE GREENWICH PHASES δ (DEG)

$$\text{TABLE 8N: } 1° \times 1° \; O_1 \; \text{OCEAN TIDE GREENWICH PHASES } \delta \; (DEG)$$

QUEEN ELIZABETH ISLANDS

GREENLAND

BAFFIN ISLAND

EASTERN CANADA

NEWFOUNDLAND

NOVA SCOTIA

EASTERN USA

LONG ISLAND

FLORIDA

BAHAMAS

CUBA

HISPANIOLA

TABLE 9N: 1° x 1° O₁ OCEAN TIDE AMPLITUDES ξ (CM)

Data table/map of ocean tide amplitudes with geographic labels: GREENLAND, ICELAND, BRITISH ISLES, FRANCE, IBERIA, MADEIRA, AZORES, NORTHWESTERN AFRICA, CANARY IS.

TABLE 9N: 1° x 1° O_1 OCEAN TIDE GREENWICH PHASES δ (DEG)

GREENLAND

ICELAND

BRITISH ISLES

IBERIA

FRANCE

NORTHWESTERN AFRICA

TABLE 1M: 1° x 1° O₁ OCEAN TIDE AMPLITUDES ξ (CM)

CENTRAL AFRICA

TABLE 1M: 1° × 1° O₁ OCEAN TIDE GREENWICH PHASES δ (DEG)

CENTRAL AFRICA

N\M	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	...	33	34	35	36	37	38	39	40	41	42
85	313	312	311	310	30.9																					
86	316	316	316	316	314																					
87	311	311	312	312	311	310																				
88	305	306	307	307	302	312	307	306	305																	
89	299	300	301	303	304	305	302	307	307	304																
90	292	294	296	298	299	299	301	303	302	303																
91	285	288	290	292	294	296	291	299	300	302																
92	278	281	284	287	289	293	294	295	300	300																
93	271	275	277	281	284	288	291	296	295	301	298															
94	264	268	272	275	278	282	286	290	293	292	297	296														
95	257	261	265	268	272	275	279	283	287	282	284	288														
96	251	255	258	262	265	268	271	275	278	282	284	282														
97	246	250	252	255	258	261	264	267	270	274	271	282														
98	241	244	247	249	252	255	257	260	263	266	265	274	274													
99	237	239	242	247	246	249	251	253	256	258	261	265	265													
100	232	235	237	239	241	243	245	247	249	251	254	257	259													
101	229	231	233	235	237	239	240	242	244	245	247	249	253	251												
102	226	228	230	231	233	234	236	237	235	240	241	243	245	249												
103	224	226	228	228	229	231	232	233	234	235	232	234	238	242												
104	220	222	223	225	226	228	229	230	231	231	232	234	235													
105	217	219	221	222	224	225	226	227	228	228	229	230														
106	215	217	218	220	221	222	223	224	225	226	226	229														
107	213	215	216	218	219	220	221	222	223	224	225	228														
108	211	213	215	216	218	219	220	221	222	223	224	227	226							360	360	360				
109	210	212	214	215	217	218	219	220	221	222	224	226	225							352	353	353	359			
110	208	210	212	213	215	217	218	220	221	222	224	226	226					337	345	340	353	354	360			
111	208	210	212	213	214	216	217	218	221	223	224	225	225					327	330	336	353	355	359			
112	207	209	211	213	215	216	218	220	221	222	223	226	224					315	326	331	356	356	355	1		
113	207	209	211	213	214	216	218	220	222	223	224	226	223							339	356	356	2	5		
114	206	208	210	212	214	216	218	220	222	223	224	226	222							342	357	351	1	7		
115	206	208	210	212	214	216	218	220	222	223	225	226	221								351	350	0	8		
116	206	208	210	213	215	216	218	220	222	224	225	226	222								340	346	353	14		
117	206	208	211	213	215	217	219	221	224	225	226	227	223								336	343	351	355		
118	206	209	212	214	216	218	220	222	225	224	227	227	224								331	339	342	356	7	

TABLE 2M: 1° x 1° O₁ OCEAN TIDE AMPLITUDES ξ (CM)

$$\text{TABLE 2M: } 1° \times 1° \; O_1 \text{ OCEAN TIDE AMPLITUDES } \xi \text{ (CM)}$$

WESTERN INDIA

MALDIVE LACCADIVE

CHAGOS

PAKISTAN

IRAN

ARABIA

CENTRAL EAST AFRICA

MADAGASCAR

TABLE 2M: 1° x 1° O₁ OCEAN TIDE GREENWICH PHASES δ (DEG)

WESTERN INDIA

PAKISTAN

IRAN

ARABIA

MALDIVE

CHANGOS

CHELL

SEY

SOC

CENTRAL EAST AFRICA

MADAGASCAR

TABLE 3M: 1° x 1° O₁ OCEAN TIDE AMPLITUDES ξ (CM)

EASTERN INDIA
SOUTHEAST ASIA
SOUTHEASTERN CHINA
NORTHWESTERN AUSTRALIA

SRI LANKA
BANGLADESH
BURMA
NICOBAR AND
THAILAND
SUMATRA
SUNDA ISLANDS
BORNEO
CELEBES
SULU SEA
PHILIPPINES
TAIWAN
HAINAN

TABLE 3M: 1° x 1° O_1 OCEAN TIDE GREENWICH PHASES δ (DEG)

EASTERN INDIA

SRI LANKA

BANGLADESH

BURMA

THAILAND

SOUTHEAST ASIA

SUMATRA

SUNDA ISLANDS

BORNEO

CELEBES

SULU SEA

PHILIPPINES

TAIWAN

HAINAN

SOUTHEASTERN CHINA

NORTHWESTERN AUSTRALIA

TABLE 4M: 1° x 1° O₁ OCEAN TIDE AMPLITUDES ξ (CM)

EASTERN CHINA

GULF OF CHIHLI

KOREA

SEA OF JAPAN

SOUTHERN JAPAN

NANPOS

MARIANAS

TAIWAN

PHILIPPINES

CELEBES

SULU SEA

SUNDA ISLANDS

CERAM SEA

NEW GUINEA

NORTHERN AUSTRALIA

BISMARCK

SOLOMON

TABLE 4M: 1° x λ_1, OCEAN TIDE GREENWICH PHASES δ (DEG)

EASTERN CHINA

GULF OF CHIHLI

SEA OF JAPAN

SOUTHERN JAPAN

KOREA

TAIWAN

NANPOS

MARIANAS

PHILIPPINES

CERAM SEA

SULU SEA

CELEBES

SUNDA ISLANDS

NEW GUINEA

BISMARCK

SOLOMON

NORTHERN AUSTRALIA

TABLE 5M: 1° x 1° O₁ OCEAN TIDE AMPLITUDES ξ (CM)

TABLE 5M: 1° x 1° O₁ OCEAN TIDE GREENWICH PHASES δ (DEG)

TABLE 6M: 1° x 1° O_1 OCEAN TIDE AMPLITUDES ξ (CM)

USA

CALIFORNIA

TABLE 6M: 1° x 1° O₁ OCEAN TIDE GREENWICH PHASES δ (DEG)

USA

CALIFORNIA

TABLE 7M: 1° x 1° O₁ OCEAN TIDE AMPLITUDES ξ (CM)

SOUTHERN USA

CALIFORNIA

MEXICO

MIDDLE AMERICA

FLORIDA

CUBA

S. AMERICA

TABLE 7M: 1° x 1° O₁ OCEAN TIDE GREENWICH PHASES δ (DEG)

SOUTHERN USA

CALIFORNIA

MEXICO

MIDDLE AMERICA

FLORIDA

CUBA

AMERICA

TABLE 8M: 1° x 1° O₁ OCEAN TIDE AMPLITUDES ξ (CM)

EASTERN USA

LONG ISLAND

FLORIDA

C2N

BAHAMAS

CUBA

JAM

HISPANIOLA

MIDDLE AMERICA

CARIBBEAN

PRO

B5A

C4

NORTHERN SOUTH AMERICA

TABLE 8M: 1° x 1° O₁ OCEAN TIDE GREENWICH PHASES δ (DEG)

$$\text{TABLE 8M: } 1° \times 1° \text{ O}_1 \text{ OCEAN TIDE GREENWICH PHASES } \delta \text{ (DEG)}$$

Region labels appearing within the table grid:

- EASTERN USA
- FLORIDA
- BAHAMAS
- CUBA
- JAM
- MIDDLE AMERICA
- LONG ISLAND
- HISPANIOLA
- PRD
- NORTHERN SOUTH AMERICA

Column headings (λ): 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320

Row index (φ): 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118

TABLE 9M: 1° x 1° O_1 OCEAN TIDE AMPLITUDE ξ (CM)

IBERIA

NORTHWESTERN AFRICA

MADEIRA

CANARYS

CAPE VERDE

AZORES

EASTERN BRAZIL

TABLE 9M: 1° x 1° O₁, OCEAN TIDE GREENWICH PHASES δ (DEG)

EASTERN BRAZIL

IBERIA

NORTHWESTERN AFRICA

λ° / φ°	319	320	321	322	323	324	325	326	327	328	329	330	331	332	333	334	335	336	337	338	339	340	341	342	343	344	345	346	347	348	349	350	351	352	353	354	355	356	357	358	359	360

TABLE 1S: 1° x 1° O$_1$ OCEAN TIDE AMPLITUDES ξ (CM)

SOUTHERN AFRICA

ANTARCTICA

TABLE 1S: 1° x 1° O₁ OCEAN TIDE GREENWICH PHASES δ (DEG)

SOUTHERN AFRICA

ANTARCTICA

N\E	357	358	359	360	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38

(Large numerical data table of O₁ ocean tide Greenwich phases in degrees; values not individually transcribable with confidence.)

TABLE 28: 1° x 1° O₁ OCEAN TIDE AMPLITUDES ξ (CM)

ANTARCTICA

TABLE 2S: 1° x 1° O_1 OCEAN TIDE GREENWICH PHASES δ (DEG)

EASTERN AFRICA

MADAGASCAR

CHAGOS

ANTARCTICA

TABLE 3S: 1° × 1° O_1 OCEAN TIDE AMPLITUDES ξ (CM)

SUNDA ISLANDS

WESTERN AUSTRALIA

ANTARCTICA

TABLE 3S: 1° x 1° O_1 OCEAN TIDE GREENWICH PHASES δ (DEG)

SUNDA ISLANDS

WESTERN AUSTRALIA

ANTARCTICA

TABLE 4S: 1° x 1° O$_1$ OCEAN TIDE AMPLITUDES ξ (CM)

SOLOMON

NEW GUINEA

ARU

TAN

SUNDA ISLANDS

CENTRAL EASTERN AUST RALIA

TASMANIA

ANTARCTICA

TABLE 4S: 1° x 1° O₁ OCEAN TIDE GREENWICH PHASES δ (DEG)

SUNDA ISLANDS

ARU

NEW GUINEA

SOLOMON

TAM

MLIW

CENTRAL EASTERN AUSTRALIA

TASMANIA

ANTARCTICA

TABLE 5S: 1° x 1° O₁ OCEAN TIDE AMPLITUDES ξ (CM)

TABLE 5S: 1° x 1° O_1 OCEAN TIDE GREENWICH PHASES δ (DEG)

Full-page numerical data table of ocean tide Greenwich phases, with geographic labels including COOK, SAMOA, TONGA, FIJI, NEW CALEDONIA, NEW HEBR, NEW ZEALAND, SOLOMON, and ANTARCTICA. Column headings run from 160 to 201 (longitude, °M) across the top; row labels run from 98 to 168 (°N) down the left side.

TABLE 6S: 1° x 1° O_1 OCEAN TIDE AMPLITUDES ξ (CM)

LATITUDE

LONGITUDE

M3E

ANTARCTICA

TABLE 6S: 1° x 1° O₁ OCEAN TIDE GREENWICH PHASES δ (DEG)

ANTARCTICA

TABLE 7S: 1° x 1° O₁ OCEAN TIDE AMPLITUDES ξ (CM)

PERU

CHILE

ANTARCTICA

ESR

TABLE 7S: 1° x 1° O₁ OCEAN TIDE GREENWICH PHASES δ (DEG)

$$\text{TABLE 7S: } 1° \times 1° \; O_1 \text{ OCEAN TIDE GREENWICH PHASES } \delta \text{ (DEG)}$$

PERU

CHILE

ANTARCTICA

TABLE 8S: 1° x 1° O_1 OCEAN TIDE AMPLITUDES ξ (CM)

SOUTHERN SOUTH AMERICA

FALKLAND

ANTARCTICA

TABLE 8S: 1° x 1° O₁ OCEAN TIDE GREENWICH PHASES δ (DEG)

SOUTHERN SOUTH AMERICA

ANTARCTICA

FALKLAND

TABLE 9S: 1° x 1° O_1 OCEAN TIDE AMPLITUDES ξ (CM)

EASTERN BRAZIL

ASCENSION

ANTARCTICA

TABLE 9S: 1° x 1° O₁ OCEAN TIDE GREENWICH PHASES δ (DEG)

*EASTERN BRAZIL*

ASCENSION

ANTARCTICA

APPENDIX B

ATLAS OF GLOBAL O$_1$ OCEAN TIDE
**CORANGE AND COTIDAL MAPS**

APPENDIX B

## ATLAS OF CORANGE AND COTIDAL MAPS
OF THE O_1 OCEAN TIDE

Amplitudes ξ of corange lines in cm.

Greenwich phases δ of cotidal lines in 15, 30, 45, 60, 75, 90, 105, 120, 135, 150, 165, 180, 195, 210, 225, 240, 255, 270, 285, 300, 315, 330, 345, 360 = 0° where 15° ≈ 1 hour.

CORANGE MAP OF O_1 OCEAN TIDE
AMPLITUDES ξ IN CM
◉ AMPHIDROMES

COTIDAL MAP OF O_1 OCEAN TIDE
GREENWICH PHASES δ IN DEGREES
15° ≈ 1 HOUR
◎ AMPHIDROMES

ARCTIC CORANGE MAP OF O₁ OCEAN TIDE

AMPLITUDES ξ IN CM

◎ AMPHIDROMES   ∗ P NORTH POLE

ARCTIC COTIDAL MAP OF O_1 OCEAN TIDE

GREENWICH PHASES δ IN DEGREES

$15° \approx 1$ HOUR

◎ **AMPHIDROMES**   * **P NORTH POLE**

DISTRIBUTION

DISTRIBUTION (Continued)

National Science Foundation
1951 Constitution Ave., N.W.
Washington, DC 20550
ATTN: Mathematical Sciences Division
       J. G. Gross
       R. E. Wall

Scripps Institution of Oceanography
University of California at San Diego
LaJolla, CA 92037
ATTN: Dr. W. H. Munk
      Dr. M. C. Hendershott
      Prof. B. D. Zetler
      Prof. S. M. Smith
      Prof. H. W. Menard
      Dr. J. H. Filloux
      Dr. B. Bernstein
      Dr. D. S. Luther

Dr. C. Wunsch
MIT/Dept. Earth & Planetary Sciences
Cambridge, MA 02139

Woods Hole Oceanographic Institute
Woods Hole, MA 02543
ATTN: Dr. H. M. Stommel
      Dr. G. Veronis
      Dr. N. P. Fofonoff
      Dr. J. Whitehead
      Dr. P. G. Brewer

Battelle Columbus Laboratories
505 King Ave.
Columbus, OH 43201
ATTN: A. G. Mourad

Dr. J. W. Chamberlain
Rice University
Houston, TX 77001

Dr. R. H. Rapp
Ohio State University
Dept. of Geodetic Science
1958 Neil Ave.
Columbus, OH 43210

Dr. R. O. Reid
Texas A&M University
College Station, TX 77843

Florida State University
Dept. of Oceanography
Tallahassee, FL 32306
ATTN: Dr. J. J. O'Brien
      Dr. W. Sturges
      Ms. L. Vasant

Prof. F. E. Snodgrass
Inst. of Geophysics and Planetary Physics
University of California at San Diego
LaJolla, CA 92037

Prof. K. Wyrtki
University of Hawaii
Honolulu, HI 96822

Prof. B. Tapley
Dept. of Aerospace Eng. & Eng. Mechanics
WRW 402
University of Texas
Austin, TX 78712

Prof. D. Lynch
Thayer School of Engineering
Dartmouth College
Hanover, NH 03755

Air Force Geodetic Laboratory
L. G. Hanscom Field
Bedford, MA 01730

David T. Haislip
U. S. Coast Guard
400 7th Street, S.W.
Washington, DC 20590

NOAA/National Ocean Survey
National Geodetic Survey
Rockville, MD 20852
ATTN: Dr. B. Chovitz
        Dr. J. M. Diamante
        Dr. B. C. Douglas
        Dr. C. C. Goad
        Dr. F. Morrison

NOAA/National Ocean Survey
Oceangraphic Division
Rockville, MD 20852
ATTN: D. C. Simpson
        D. L. Porter
        R. A. Smith
        B. B. Parker

NOAA Atlantic Oceanographic and
  Meteorological Lab.
Physical Oceanography Laboratory
15 Rickenbacker Causeway
Miami, FL 33149
ATTN: G. A. Maul
        H. M. Byrne

NOAA Pacific Marine Environmental Lab.
Seattle, WA 98105
ATTN: Dr. J. R. Apel
        H. O. Mofield
        C. A. Pearson
        J. Byrne

NOAA/Geophysical Fluid Dynamics Lab.
Princeton University
Princeton, NJ 08540
ATTN: Dr. J. Smagorinsky
        Dr. K. Bryan
        Dr. M. D. Cox

NOAA/National Center for Atmospheric
  Research
Boulder, CO 80303
ATTN: Dr. W. R. Holland

NASA/Goddard Space Flight Center
Greenbelt, MD 20771
ATTN: Dr. J. W. Siry
        D. E. Smith
        J. G. Marsh
        T. L. Felsentreger
        J. Zwally

NASA/Wallops Station
Information Processing and Analysis
  Branch
Wallops Island, VA 23337
ATTN: C. D. Leitao
        N. E. Huang
        W. B. Krabill
        B. Speidel

Director
U.S. Army Ballistic Research Laboratory
Aberdeen Proving Ground, MD 21005
ATTN: DRDAR-TBS-S (STINFO)

Smithsonian Astrophysical Observatory
60 Garden St.
Cambridge, MA 02138
ATTN: Dr. E. M. Gaposchkin
        Dr. G. C. Wiffenbach
        B. Stevens

DISTRIBUTION (Continued)

University of Washington
Dept. of Oceanography WB-10
Seattle, WA 98195
ATTN: Dr. A. J. Clarke
        Prof. D. Winter
        Prof. M. Jamart

Jet Propulsion Laboratory
California Institute of Technology
4800 Oak Grove Drive
Pasadena, CA 91103
ATTN: Dr. M. Parke
        Dr. J. Lorell
        Dr. G. Born

Prof. J. T. Kuo
Lamont-Doherty Geological Observatory
Columbia University
Palisades, NY 10964

Prof. W. J. Pierson, Jr.
1641 Rosalind Ave.
Elmont, NY 11003

Prof. Morris Schulkin
Applied Physics Laboratory
University of Washington
Seattle, WA 98105

Dr. R. H. Estes
Business and Technological Systems, Inc.
Aerospace Building, Suite 605
10210 Greenbelt Rd.
Seabrook, MD 20801

T. V. Martin
Sci. Res. and Appl. Group
Washington Analytical Services Center, Inc.
6801 Kenilworth Ave.
Riverdale, MD 20840

Dr. S. K. Jordan
The Analytic Sciences Corporation
6 Jacob Way
Reading, MA 01867

The Rand Corporation
Santa Monica, CA 90406
ATTN: Director, Climate Program

Local:

C
D
E31 (GIDEP)
E41
K
K02
K04
K05
K10
K102
K1040
K104U
K104S   (300)
K104Z
K12
K12G
K12M
K12S
K12T
K12 W
K13
K13H
K14
K14S
K20
K204
K21
K21D
K30

K40
K404A
K404S
K41
K42
K44
K50
K51B
R
R02
R04
R31G
R40
R44
R44VT
X210   (6)